BMC Endocrine Disorders

Open Access

Validation of the Persian version of the summary of diabetes self-care activities scale (SDSCA) in pregnant women with gestational diabetes mellitus using a COSMIN methodology



Mahsa Maghalian¹, Mohammad Asghari Jafarabadi^{2,3,4}, Fatemeh Abbasalizadeh⁵, Sakineh Mohammad-Alizadeh-Charandabi¹, Solmaz Ghanbari-Homaie¹ and Mojgan Mirghafourvand^{6*}

Abstract

Background Gestational diabetes mellitus (GDM) is a condition with significant prenatal and postnatal implications. This study aimed to validate the Summary of Diabetes Self-Care Activities (SDSCA) measure in Iranian women with GDM, focusing on its psychometric properties.

Methods The Persian version of the SDSCA was evaluated in 180 Iranian women with GDM. Following COSMIN guidelines, the instrument was translated into Persian, and its psychometric properties were assessed, including content validity, face validity, construct validity, internal consistency, test-retest reliability, measurement error, responsiveness, and interpretability. Floor and ceiling effects were also examined.

Results The validity assessments showed strong content validity, with a Content Validity Index (CVI) of 0.93 and a Content Validity Ratio (CVR) of 0.97. Face validity yielded an impact score of 4.38. Exploratory factor analysis (EFA) identified three factors—diet, exercise, and blood sugar testing—accounting for 57.4% of the variance. Confirmatory factor analysis (CFA) confirmed the model's excellent fit (CFI = 1.00, TLI = 0.99, NFI = 0.98, RFI = 0.96). The reliability analysis showed a Cronbach's alpha of 0.78 and a McDonald's omega of 0.91, with an intraclass correlation coefficient (ICC) of 0.92 (95% CI: 0.83–0.96). Ceiling effects were observed for blood sugar testing (26.7%), while floor effects were noted for exercise (6.7%) and blood sugar testing (6.1%). The Minimal Important Change (MIC) of 2.68 units exceeded the Smallest Detectable Change (SDC) of 1.11 units, indicating the tool's ability to detect clinically meaningful changes.

Conclusions The Persian version of the SDSCA demonstrates strong psychometric properties, including both reliability and validity, making it a suitable tool for assessing self-care behaviors in Iranian women with GDM. Its use in future research can enhance understanding of self-management in this population.

*Correspondence: Mojgan Mirghafourvand mirghafourvand@gmail.com

Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by-nc-nd/4.0/.

Keywords Cross-Sectional studies, Self care, Psychometrics, Pregnancy, Diabetes, Gestational, Iran

Background

Diabetes is a metabolic disease marked by challenges in regulating blood sugar levels due to insufficient insulin production, insulin resistance, or both. It poses a significant public health issue, with projections indicating that nearly 1 in 10 adults globally will be affected by 2035 [1, 2]. Uncontrolled diabetes can lead to severe complications, including cardiovascular diseases, cognitive decline, and depression [3]. The condition is classified into several types, namely Type 1, Type 2, and gestational diabetes mellitus (GDM) [4, 5].

GDM, defined as glucose intolerance during pregnancy, affects approximately 14% of pregnancies worldwide, with its prevalence rising alongside obesity and Type 2 diabetes rates [6]. It is more widespread than previously recognized, impacting women across various body weights and regions, including Asia and Europe. In Iran, the prevalence of diabetes is around 7.6% in the general population [7].

Key risk factors for GDM include family history of diabetes, obesity, previous GDM, multiparity, and miscarriage history. Modifiable factors such as unhealthy diets and physical inactivity also play a role, with women from racial or ethnic minority groups at heightened risk [8, 9].

GDM is linked to negative outcomes for both mothers and their offspring, including preterm birth, cesarean delivery, and long-term metabolic risks for the child. Women with GDM are also more likely to develop Type 2 diabetes and cardiovascular diseases later in life [10, 11]. This condition imposes a substantial burden on health-care systems due to the need for enhanced monitoring and management during and after pregnancy [12].

Management of GDM focuses on self-care practices like blood glucose monitoring, dietary changes, and increased physical activity [13]. Empowerment-based interventions have proven effective in improving self-care behaviors and maternal health outcomes [14]. The World Health Organization (WHO) emphasizes the importance of self-care in health maintenance [15]. Adherence to self-care is vital for effective diabetes management and glycemic control [16]. A personalized, patient-centered approach is essential for helping women maintain healthy behaviors and prevent future diabetes onset [17, 18].

Despite the importance of self-care, standardized tools to assess these behaviors in women with GDM are lacking. While instruments like the Summary of Diabetes Self-Care Activities (SDSCA) have been validated for general diabetes populations [19–22], their suitability for GDM, particularly in diverse cultural contexts, remains underexplored. Recent studies have validated the SDSCA in Hindi [23] and Arabic [24] for women with GDM, highlighting its potential utility.

Utilizing validated instruments to measure healthrelated patient-reported outcomes (HR-PROs) is crucial for effective disease management [25]. The COSMIN (Consensus-Based Standards for the Selection of Health Status Measurement Instruments) checklist offers a framework for evaluating the psychometric properties of such instruments [25, 26]. Given the increasing prevalence of GDM and its complications, there is a pressing need for validated tools to assess self-care behaviors in specific populations. This study aims to psychometrically validate the Persian version of the SDSCA among Iranian women with GDM, following COSMIN guidelines to evaluate the scale's validity, reliability, and other psychometric properties for the target population.

Methods

Study participants and setting

Before using the SDSCA, the researchers obtained the necessary permissions from the SDSCA working group [22]. The study was approved by the Ethics Committee of Tabriz University of Medical Sciences in Iran. The validation was conducted with a sample of 180 pregnant women diagnosed with GDM. The participants were recruited from the outpatient clinics of three government-operated hospitals in Tabriz city, namely, Taleghani, Al-Zahra, and 29 Bahman, between February 23, 2024, and July 12, 2024. The sample was divided into two groups: 80 participants for exploratory factor analysis and 100 participants for confirmatory factor analysis. All women provided written informed consent before participating in the study.

The eligibility criteria included a GDM diagnosis, age \geq 18 years, and gestational age \geq 24 weeks. GDM was diagnosed if one or more of the following plasma glucose values were met or exceeded during a 75-gram oral glucose tolerance test (OGTT): fasting glucose \geq 92 mg/dL (5.1 mmol/L), 1-hour glucose \geq 180 mg/dL (10.0 mmol/L), or 2-hour glucose \geq 153 mg/dL (8.5 mmol/L) [27]. The exclusion criteria were preexisting diabetes, fetal abnormalities, severe medical conditions, and maternal psychiatric disorders.

Instruments

Sociodemographic and obstetrics checklist

The sociodemographic and obstetric checklist collected information on the participants' age, occupation, education, family income, and obstetric history, including number of pregnancies and parity.

Summary of the diabetes Self-Care activities (SDSCA)

The SDSCA instrument was originally developed in the United States by Toobert and colleagues in 2000. It is a comprehensive 11-item measure that assesses various aspects of diabetes self-management, including dietary practices (4 items), exercise habits (2 items), blood glucose monitoring (2 items), foot care (2 items), and smoking behavior (1 item). Each item asks participants to report the number of days in the past week they engaged in a specific self-care behavior. Responses are recorded on a 7-point scale, ranging from 0 (no days) to 7 (every day). For this study [22], the Persian version of the SDSCA was adapted to include 7 items focusing on three key domains relevant to GDM: diet (3 items), exercise (2 items), and blood glucose testing (2 items). We excluded foot care and smoking from the SDSCA as they are less relevant to GDM management. Foot care primarily concerns those with chronic diabetes, while smoking's prevalence among pregnant women in Iran is low. Instead, we focused on key self-care behaviors like diet, exercise, and blood glucose monitoring that are more pertinent to GDM.

Sample size determination

The study adhered to recommended guidelines for factor analysis sample size [28], initially targeting 80 participants for exploratory factor analysis (EFA) based on 8 survey items. To ensure robust validation through both EFA and confirmatory factor analysis (CFA) on separate datasets, the total sample size was increased to 180 participants, with 80 allocated for EFA and 100 for CFA [28, 29]. The CFA sample size was determined by the rule of having at least 10 participants per free parameter, with 7 free parameters requiring a minimum of 70 participants [29]. The inclusion of 100 participants for CFA exceeded this requirement, ensuring sufficient statistical power.

Statistical analyses

Statistical analyses were performed using IBM SPSS Statistics 22 and STATA 14. Continuous variables with normal distributions were reported as means and standard deviations, while categorical variables were presented as frequencies and percentages.

Translation procedure

The SDSCA questionnaire was translated into Persian following WHO guidelines. This involved forward translation by two Persian speakers, back-translation by two native English speakers, and reconciliation of discrepancies [30]. A pilot study with 10 eligible women assessed the comprehensibility and ease of use of the Persian version, with feedback incorporated into the final questionnaire [31].

Validity assessment

Content validity was assessed by 10 experts using content validity index (CVI > 0.79) and content validity ratio (CVR > 0.62) [32].

Face validity was evaluated by a separate group of 10 eligible women, with an impact score > 1.5 considered acceptable [33].

Construct validity was examined through EFA and CFA. We used both EFA and CFA following the twostep procedure outlined by Malik and Millsap [34, 35]. EFA was employed to explore the underlying factor structure and identify the appropriate number of factors, while CFA was used to confirm the significance of the relationships between the factors and observed variables, ensuring model fit and validity. EFA utilized principal axis factoring with promax rotation, and the Kaiser-Meyer-Olkin (KMO) measure was greater than 0.5, along with Bartlett's test of sphericity. The minimum cut-off point for factor loadings in the EFA was set at 0.3 [35]. CFA assessed model fit using the following criteria: RMSEA < 0.08, SRMR < 0.10, normed chi-square (χ^2 / df) < 5, and comparative fit indices (CFI, NFI, RFI, TLI, GFI>0.90) [35, 36].

Reliability assessment

Internal consistency was evaluated using Cronbach's alpha and McDonald's omega, with values ≥ 0.7 considered acceptable for all 180 participants [28, 37].

Test-retest reliability was assessed by administering the questionnaire to 30 participants twice, 14 days apart. Intraclass correlation coefficient (ICC) values>0.7 indicated good reliability [38].

Additional analyses

Floor and ceiling effects were considered significant if > 15% of responses fell at the extremes [39].

Responsiveness was assessed by comparing SDC to the minimal important change (MIC), with SDC < MIC indicating adequate responsiveness [26].

Interpretability was evaluated by estimating MIC as half the standard deviation of instrument scores [40, 41].

Measurement error was evaluated using the standard error of measurement (SEM) and smallest detectable change (SDC), with lower SDC values indicating higher sensitivity [42].

Results

Descriptive characteristics of the participants

The researchers approached 207 women for participation in the validation study. Of these, 11 were excluded because they did not meet the eligibility criteria. Among the remaining 196 eligible participants, 16 declined to participate, resulting in a final sample of 180 women and a response rate of 91.8%. The study included a total of 180

Table 1	Demographic characteristics of the participants in	the
SDSCA (r	n = 180)	

Characteristics	EFA (n = 8	80)	CFA (n = 100)		
	Mean	SD	Mean	SD	
Age (year)	33.06	6.91	31.63	7.30	
Spouse age (year)	38.3	7.22	36.41	7.13	
Parity	0.98	0.86	0.96	0.92	
Gravity	2.53	1.33	2.36	1.24	
	Number	Percent	Number	Percent	
Education level					
High school or below	39	48.9	46.0	46.0	
Diploma and university	41	51.1	54.0	54.0	
Job					
Housewife	76	95	83.0	83.0	
Employee	4	5	17.0	17.0	
Family history of diabetes	33	41.3	30.0	30.0	
Income sufficiency					
Insufficient	15	8.3	13	13	
Relatively sufficient	69	86.3	84	84	
Completely sufficient	2	2.5	3	3	

SD=standard deviation, EFA=exploratory factor analysis, CFA=confirmatory factor analysis, SDSCA=summary of diabetes self-care activities scale

 Table 2
 Results for the content and face validity of the SDSCA

Items	Impact score	CVI	CVR
1	5	0.90	1
2	3.70	0.86	1
3	3.60	0.93	1
4	4.10	0.90	1
5	4.60	1	1
6	5	1	1
7	4.70	0.90	0.80
Total	4.38	0.93	0.97

CVI=content validity index, CVR=content validity ratio, SDSCA=Summary of Diabetes Self-Care Activities Scale

women, who were randomly divided into two groups: an EFA group of 80 participants and a CFA group of 100 participants. The characteristics of the participants are detailed in Table 1. The women in the EFA group had a mean age of 33.1 years, with a SD of 6.9 years. In the CFA group, the women had a mean age of 31.6 years, with a SD of 7.3 years.

The majority of women in both groups were housewives, comprising 95% of the EFA group and 83% of the CFA group. Additionally, most participants reported relatively sufficient incomes, with 86.3% in the EFA group and 84% in the CFA group indicating income adequacy.

The mean (SD) scores for each factor were as follows: diet, 4.55 (1.55); exercise, 3.67 (2.10); and blood sugar test, 4.52 (2.33). For the overall SDSCA tool, the mean (SD) score was 4.29 (1.42).

Validity assessment

Content validity The CVI was 0.93 and the CVR was 0.97, indicating strong content validity (Table 2).

Face validity The impact score was 4.38, indicating strong face validity (Table 2).

Construct validity During the EFA process, one item (item 4) was excluded because its factor loading was less than 0.3 [35], ultimately reducing the number of items from 8 to 7 (KMO = 0.64, Bartlett's test of sphericity p < 0.001). The results presented in Table 3 show the extracted factors along with the corresponding questionnaire items. The first factor was labeled "diet." This factor contains 3 items and explains 20.2% of the total variance. The second factor, "exercise," consisted of 2 items and accounted for 17.5% of the overall variance. The third factor was "blood sugar testing." This factor also had 2 associated items and made up 19.7% of the total variance (Figs. 1 and 2).

We conducted a CFA to evaluate the three-factor structure from the previous EFA. The analysis revealed a chisquare to degrees of freedom ratio (χ^2 /df) of 1.25 (χ^2 = 13.69, df=11), which falls within the acceptable range. This finding indicates that the model has a good fit with the data. Additionally, the key fit indices, including the TLI, CFI, NFI, GFI, and RFI, all exceeded the recommended threshold of 0.9, further confirming the overall goodness of fit of the model. Importantly, the RMSEA value of 0.04 and the SRMR value of 0.02 suggest that the model is valid and reliable, as these values are within the recommended thresholds for a well-fitting model (Table 4).

Reliability assessment

Internal consistency The reliability analysis revealed that the questionnaire had a Cronbach's alpha of 0.78 and a McDonald's omega of 0.91, indicating adequate internal consistency (Table 5).

Test-Retest Reliability: The ICC was estimated to be 0.92 (95% CI: 0.83–0.96) (Table 5).

Additional analyses

Floor and ceiling effects The ceiling effect on the overall SDSCA score was 0.6%. When the individual subdomains were examined, the ceiling effects were found to be 1.1% for the Diet component, 3.3% for Exercise, and 26.7% for the Test of Blood Sugar subdomain. Additionally, the analysis revealed a floor effect of 0.6% on the overall SDSCA score, whereas the specific subdomains presented floor effects of 1.1% for diet, 6.7% for exercise, and 6.1% for the test of the blood sugar component (Table 5).

Responsiveness and interpretability The MIC value of 2.68 units was greater than the SDC value of 1.11 units,

Table 3 Fracture structure of the SDSCA

Scale item	Fac- tor 1	Fac- tor 2	Fac- tor 3
Factor1: Diet			
How many of the last seven days have you followed a healthful eating plan?	0.84		
On average, over the past month, how many days per week have you followed you're eating plan?	0.83		
On how many of the last seven days did you eat five or more servings of fruits and vegetables?	0.40		
On how many of the last seven days did you eat high fat foods such as red meat or full-fat dairy products?	0.07		
Factor2: Exercise			
On how many of the last seven days did you participate in at least 30 min of physical activity? (Total minutes of continuous activ- ity, including walking).		0.83	
On how many of the last seven days did you participate in a specific exercise session (such as swimming, walking, biking) other than what you do around the house or as part of your work?		0.83	
Factor3: Blood Sugar Testing			
On how many of the last seven days did you test your blood sugar?			0.95
On how many of the last seven days did you test your blood sugar the number of times recommended by your health care provider?			0.95
% of variance observed	20.2	17.5	19.7
Total score	57.4		

SDSCA = Summary of diabetes self-care activities scale



Fig. 1 Factor structure model plot of the SDCSCA based on CFA Dit = diet, Exr = exercise, and BST = test of blood sugar

indicating that the measurement tool is able to accurately identify clinically or practically meaningful changes in the parameter being measured. This suggests that the tool has sufficient responsiveness and is able to consistently detect meaningful changes (Table 5).

Measurement error The SEM was calculated as 0.40, meaning that the recorded values are expected to fall

within ± 0.40 units of the true score. The SDC was determined to be 1.11 units, indicating that any variation in the measured quantity less than this threshold may be imperceptible due to measurement uncertainties and can therefore be regarded as insignificant (Table 5).



Fig. 2 Factor load scree plot of the items for determining the number of extracted factors of the Persian version of SDSCA (summary of diabetes self-care activities scale)

Table 4 N	lodel fit indicato	rs of the SDSCA
-----------	--------------------	-----------------

Goodness of fit indices	Value
	13.69
df	11
Chi2/df	1.25
P value	0.25
CFI	1
TLI	0.99
NFI	0.98
RFI	0.96
GFI	1.00
SRMR	0.02
RMSEA	0.04

Df=degrees of freedom, χ^2/df =normed chi-square, GFI=goodness-of-fit index, RFI=relative fit index, NFI=normed fit index, CFI=comparative fit index, TLI=Tucker–Lewis index, SRMR=standardized root mean square residual, RMSEA=root mean square error of approximation, SDSCA=summary of diabetes self-care activities scale

Discussion

This study is the first to evaluate the measurement properties of the SDSCA instrument in Iranian women with GDM, following the COSMIN checklist. The results confirm the validity, reliability, responsiveness, and interpretability of the SDSCA tool within this population.

Effective self-care behaviors are essential for managing GDM, encompassing diet, exercise, medication adherence, and glucose monitoring. The role of healthcare providers in promoting these behaviors is critical to preventing complications associated with diabetes [43, 44]. Adherence to self-care practices allows women to exert greater control over their diet and body weight [45]. The self-care practices essential for managing GDM are largely analogous to those required for type 2 diabetes management. This similarity suggests that the components of the SDSCA scale, originally designed for type 2 diabetes, can be effectively adapted to evaluate self-care behaviors in pregnant women with GDM [46]. Despite the existence of self-care assessment tools for type 2 diabetes [47-54], there are currently no validated instruments tailored for Iranian women with GDM.

The CVI of 0.93 and CVR of 0.97 indicate that the SDSCA has excellent content validity for Iranian women with GDM. These findings align with previous studies by Al Hashmi et al. [24] and Singh et al. [23], which reported CVI values between 0.8 and 1 for the same tool in other populations. The high CVI and CVR values suggest that the tool's items are highly relevant and representative of the self-care challenges faced by women with GDM [28, 55]. Additionally, the Cronbach's alpha coefficient of

Factor	Cronbach's α coefficient	McDonald's omega	ICC	(95% CI)	SEM	SDC	MIC	AVE	Floor ef- fect (%)	Ceil- ing effect
										(%)
Diet	0.71	0.74	0.96	(0.92,0.98)	0.31	0.86	2.07	0.53	1.1	1.1
Exercise	0.81	0.81	0.95	(0.89,0.97)	0.46	1.27	3.08	0.69	6.7	3.3
Blood Sugar	0.94	0.95	0.85	(0.69, 0.93)	0.90	2.49	6.03	0.90	6.1	26.7
Testing										
Total	0.78	0.91	0.92	(0.83,0.96)	0.40	1.11	2.68	0.70	0.6	0.6
					-		,			

Table 5 Stability coefficients and interclass correlation coefficient SDS
--

ICC=intraclass correlation coefficient, CI=confidence interval, SEM=standard error of measurement (SEM=SD $\sqrt{1-ICC}$), SDC=smallest detectable change (SDC=SEM1.96 $\sqrt{2}$), MIC=minimal important change (SEM*sqrt(n)/2), AVE=average variance extracted (acceptable if AVE>0.5, the threshold is 0.36–0.5), SDSCA=Summary of Diabetes Self-Care Activities Scale

0.78 demonstrates good internal consistency, consistent with the Hindi (α =0.82) and Arabic (α =0.74) versions of the tool [23, 24]. These results collectively support the robustness of the SDSCA across diverse cultural contexts.

The EFA revealed a three-factor structure, consistent with the primary self-care behaviors relevant to GDM management: diet, exercise, and blood glucose monitoring. Notably, Item 4 was excluded due to a low factor loading (< 0.3). This finding contrasts with studies by Singh et al. [23] and Al Hashmi et al. [24], where Item 4 performed adequately. However, similar issues with Item 4 have been reported in the German and Korean versions of the SDSCA for type 2 diabetes [48, 55], suggesting that its poor performance may not be unique to this study. One possible explanation is cultural differences in the interpretation of the item's content, which may not resonate equally across populations. Future research should explore the cultural and contextual factors influencing item performance to enhance the tool's cross-cultural applicability.

The factor structure identified in this study aligns with previous research on GDM self-care behaviors. For instance, diet emerged as the first factor, reflecting its central role in GDM management. Women with GDM often struggle with dietary control and require tailored recommendations from clinical nutritionists to manage carbohydrate intake and postprandial blood glucose levels [56, 57]. Exercise, the second factor, is equally critical, as physical activity improves glucose uptake by skeletal muscles and enhances glycemic control [58, 59]. However, adherence to exercise recommendations remains suboptimal among women with GDM [60]. Blood glucose monitoring, the third factor, is essential for preventing adverse pregnancy outcomes, yet adherence to self-monitoring practices is often inconsistent [51]. These findings underscore the importance of patient education and active involvement in self-care to improve outcomes and reduce long-term diabetes risk [61, 62].

The CFA results demonstrated an excellent fit for the proposed factor structure, with all key fit indices meeting

or exceeding recommended thresholds. This represents a significant advancement over previous studies [23, 24], which did not conduct CFA. The rigorous evaluation of structural validity through CFA strengthens the evidence supporting the SDSCA's use in GDM populations.

Strengths and limitations

Our study has several strengths, including adherence to the COSMIN checklist, a comprehensive evaluation of psychometric properties, and the use of a relatively large and diverse sample of Iranian women with GDM. However, we acknowledge certain limitations. First, the lack of a gold standard measure for assessing criterion-related validity limits our ability to compare the SDSCA with an established benchmark. Second, the cross-sectional design precludes the assessment of the tool's responsiveness to changes in self-care behaviors over time. Third, the potential for response bias due to self-report measures cannot be ruled out. Finally, the cross-cultural validity of the Persian version was not examined, which should be addressed in future research.

Future research directions

Future studies should focus on several key areas to build on our findings. First, longitudinal studies are needed to assess the SDSCA's responsiveness to changes in self-care behaviors over time, particularly in response to interventions aimed at improving GDM management. Second, qualitative research could provide deeper insights into the cultural and contextual factors influencing self-care practices in women with GDM, helping to refine the SDSCA for use in diverse populations. Third, the cross-cultural validity of the Persian version should be evaluated in other populations to ensure its broader applicability. Finally, future research should explore the factors contributing to the poor performance of Item 4 and consider modifying or replacing it with culturally relevant items.

Clinical implications

The Persian version of the SDSCA demonstrates promising psychometric properties for assessing self-care behaviors in Iranian women with GDM. However, the clinical adoption of this scale should be approached with caution. The lack of evaluation of cross-cultural validity and the potential limitations in cultural adaptation highlight the need for further research before the scale can be confidently implemented in clinical practice. Future studies should rigorously assess the cultural appropriateness of the scale and its applicability to diverse populations. Until then, healthcare providers should consider these limitations when interpreting the results and use the scale as a supplementary tool rather than a definitive measure of self-care behaviors in this population.

Conclusion

The Persian version of the SDSCA demonstrates strong psychometric properties, making it a suitable tool for assessing self-care behaviors in Iranian women with GDM. Its use in clinical and research settings can enhance understanding of self-management behaviors and inform targeted interventions to improve outcomes for women with GDM and their children. Given the significant consequences of uncontrolled GDM, this tool represents a valuable contribution to diabetes care in Iran.

Abbreviations

Confirmatory factor analysis
Comparative fit index
Consensus-Based Standards for the Selection of Health Status
Measurement Instruments
Content validity index
content validity ratio
Degrees of freedom
Exploratory factor analysis
Gestational diabetes mellitus
Goodness-of-fit index
Health-related patient-reported outcomes
Intraclass correlation coefficient
Kaiser–Meyer–Olkin
Minimal important change
Normed fit index
Relative fit index
Root mean square error of approximation
Standard deviation
Smallest detectable change
Summary of diabetes self-care activities scale
Standard error of measurement
Standardized root mean squared residual
Tucker–Lewis index
World Health Organization
Normed chi-square test

Acknowledgements

We would like to express our sincere gratitude to the Research Development Unit of Al-Zahra Educational and Treatment Center and the Clinical Research Development Unit of Taleghani Hospital, Tabriz University of Medical Sciences, for their invaluable scientific support, which made this research possible.

Author contributions

The study design was developed collaboratively by MMi, MMa, SMAC, SGH, and FA. The initial version of the manuscript was drafted by MMi, MMa, and SMAC, while the data analysis was conducted by MAJ. All authors carefully reviewed the text, provided feedback and revisions, and approved the final submitted manuscript.

Funding

This study was funded by Tabriz University of Medical Sciences (Plan Number: IRCT20120718010324N80; Approval Date: January 3, 2024). The funding source did not participate in the study design, execution, or publication decisions.

Data availability

The study data are not publicly available due to patient privacy and ethical restrictions but can be requested from the corresponding author.

Declarations

Ethics approval and consent to participate

The study received ethical approval (IR.TBZMED.REC.1402.652/2023-12-04), and participant consent was obtained. The research methods adhered to relevant guidelines and regulations.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Clinical trial number Not applicable.

Author details

¹Department of Midwifery, Faculty of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, Iran

²Cabrini Research, Cabrini Health, Melbourne, VIC 3144, Australia ³School of Public Health and Preventive Medicine, Faculty of Medicine, Nursing and Health Sciences, Monash University, VIC 3800, Melbourne, Australia

⁴School of Clinical Sciences, Faculty of Medicine, Nursing and Health Sciences, Monash University, Clayton, VIC 3168, Australia
⁵Women's Reproductive Health Research Center, Department of Perinatology, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

⁶Social determinants of Health Research Center, Faculty of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, Iran

Received: 5 November 2024 / Accepted: 9 April 2025 Published online: 18 April 2025

References

- Zimmet P, Alberti KG, Shaw J. Global and societal implications of the diabetes epidemic. Nature. 2001;414(6865):782–7. https://doi.org/10.1038/414782a.
- Paneni F, Cosentino F, Paneni F, Cosentino F. Epidemiology, definition, and diagnosis of diabetes mellitus. In.: Springer; 2015.
- Tomic D, Shaw JE, Magliano DJ. The burden and risks of emerging complications of diabetes mellitus. Nat Rev Endocrinol. 2022;18(9):525–39. https://doi. org/10.1038/s41574-022-00690-7.
- Pleus S, Tytko A, Landgraf R, Heinemann L, Werner C, Müller-Wieland D, et al. Definition, classification, diagnosis and differential diagnosis of diabetes mellitus: update 2023. Exp Clin Endocrinol Diabetes. 2024;132(3):112–24. http s://doi.org/10.1055/a-2166-6643.
- Ramachandran A, Snehalatha C, Raghavan A, Nanditha A. Classification and diagnosis of diabetes. *Textbook of diabetes* 2024:22–7.
- Wang H, Li N, Chivese T, Werfalli M, Sun H, Yuen L, IDF Diabetes Atlas Committee Hyperglycemia in Pregnancy Special Interest Group. IDF Diabetes Atlas, et al. Estimation of global and regional gestational diabetes mellitus prevalence for 2021 by international association of diabetes in pregnancy study group's

criteria. Diabetes Res Clin Pract. 2022;183:109050. https://doi.org/10.1016/j.di abres.2021.109050.

- Gitlin ES, Demetres M, Vaidyanathan A, Palmer N, Lee H, Loureiro S, et al. The prevalence of gestational diabetes among underweight and normal weight women worldwide: a scoping review. Front Clin Diabetes Healthc. 2024;5:1415069. https://doi.org/10.3389/fcdhc.2024.1415069.
- Semnani-Azad Z, Gaillard R, Hughes AE, Boyle KE, Tobias DK, ADA/EASD PMDI, Perng W. Precision stratification of prognostic risk factors associated with outcomes in gestational diabetes mellitus: a systematic review. Commun Med (Lond). 2024;4(1):9. https://doi.org/10.1038/s43856-023-00427-1.
- Dewi RS, Isfandiari MA, Martini S, Yi-Li C. Prevalence and risk factors for gestational diabetes mellitus in Asia: a review. J Public Health Afr. 2023;14(2):7.
- Oros Ruiz M, Perejón López D, Serna Arnaiz C, Siscart Viladegut J, Àngel Baldó J, Sol J. Maternal and fetal complications of pregestational and gestational diabetes: a descriptive, retrospective cohort study. Sci Rep. 2024;14(1):9017. h ttps://doi.org/10.1038/s41598-024-59465-x.
- 11. Ye W, Luo C, Huang J, Li C, Liu Z, Liu F. Gestational diabetes mellitus and adverse pregnancy outcomes: systematic review and meta-analysis. BMJ. 2022;377:e067946. https://doi.org/10.1136/bmj-2021-067946.
- Meregaglia M, Dainelli L, Banks H, Benedetto C, Detzel P, Fattore G. The short-term economic burden of gestational diabetes mellitus in Italy. BMC Pregnancy Childbirth. 2018;18(1):58. https://doi.org/10.1186/s12884-018-168 9-1.
- Hoffman L, Nolan C, Wilson JD, Oats JJ, Simmons D. Gestational diabetes mellitus–management guidelines. The Australasian diabetes in pregnancy society. Med J Aust. 1998;169(2):93–7. https://doi.org/10.5694/j.1326-5377. 1998;tb140192. x.
- Rahimi M, GhorbaniNia R, Azizi Y, Ovaisi Z, Mohammadian S, Aminipour F. Self-Care behaviors in pregnant women with gestational diabetes: Self-care behaviors in gestational diabetes. Galen Med J. 2024;13:e3235–3235.
- 15. World Health Organization. WHO guideline on self-care interventions for health and well-being, 2022 revision; 2022.
- Ahmad F, Joshi SH. Self-Care practices and their role in the control of diabetes: A narrative review. Cureus. 2023;15(7):e41409. https://doi.org/10.7759/cur eus.41409.
- Nankervis A, Price S, Conn J. Gestational diabetes mellitus: A pragmatic approach to diagnosis and management. Aust J Gen Pract. 2018;47(7):445–9. https://doi.org/10.31128/AJGP-01-18-4479.
- Roesler A, Butten K, Taylor P, Morrison M, Varnfield M, Holmes-Truscott E. The experiences of individuals who have had gestational diabetes: A qualitative exploration. Diabet Med. 2024;e15374. https://doi.org/10.1111/dme.15374.
- Ausili D, Barbaranelli C, Rossi E, Rebora P, Fabrizi D, Coghi C, Luciani M, Vellone E, Di Mauro S, Riegel B. Development and psychometric testing of a theorybased tool to measure self-care in diabetes patients: the Self-Care of diabetes inventory. BMC Endocr Disord. 2017;17(1):66. https://doi.org/10.1186/s1290 2-017-0218-y.
- Wang RH, Lin LY, Cheng CP, Hsu MT, Kao CC. The psychometric testing of the diabetes health promotion self-care scale. J Nurs Res. 2012;20(2):122–30. http s://doi.org/10.1097/jnr.0b013e318254eb47.
- García AA. The diabetes symptom Self-Care inventory: development and psychometric testing with Mexican Americans. J Pain Symptom Manage. 2011;41(4):715–27. https://doi.org/10.1016/j.jpainsymman.2010.06.018.
- Toobert DJ, Hampson SE, Glasgow RE. The summary of diabetes self-care activities measure: results from 7 studies and a revised scale. Diabetes Care. 2000;23(7):943–50. https://doi.org/10.2337/diacare.23.7.943.
- Mallicka, Singh SK, Pandey A, Manar MK, Srivastava S. Psychometric validation of the Hindi version of summary of diabetes self-care activities (H-SDSCA) among pregnant women with gestational diabetes mellitus, in Lucknow. J Family Med Prim Care. 2023;12(2):360–5. https://doi.org/10.4103/jfmpc.jfmpc _1935_22.
- Al Hashmi I, Al-Noumani H, Alaloul F, Murthi S, Khalaf A. Translation and psychometric validation of the Arabic version of summary of the diabetes Self-Care activities (SDSCA) among pregnant women with gestational diabetes. BMC Pregnancy Childbirth. 2022;22(1):563. https://doi.org/10.1186/s1288 4-022-04897-4.
- Mokkink LB, Prinsen CA, Bouter LM, Vet HC, Terwee CB. The COnsensus-based standards for the selection of health measurement instruments (COSMIN) and how to select an outcome measurement instrument. Braz J Phys Ther. 2016;20(2):105–13. https://doi.org/10.1590/bjpt-rbf.2014.0143.
- Mokkink LB, Terwee CB, Knol DL, Stratford PW, Alonso J, Patrick DL, Bouter LM, de Vet HC. The COSMIN checklist for evaluating the methodological quality

of studies on measurement properties: a clarification of its content. BMC Med Res Methodol. 2010;10:22. https://doi.org/10.1186/1471-2288-10-22.

- 27. International Association of Diabetes and Pregnancy Study Groups Consensus Panel, Metzger BE, Gabbe SG, Persson B, Buchanan TA, Catalano PA, Damm P, et al. International association of diabetes and pregnancy study groups recommendations on the diagnosis and classification of hyperglycemia in pregnancy. Diabetes Care. 2010;33(3):676–82. https://doi.org/10.2337/ dc09-1848.
- Hoyle RH. Confirmatory factor analysis. In: Tinsley HEA, Brown SD, editors. Handbook of applied multivariate statistics and mathematical modeling. San Diego, CA: Academic; 2000. pp. 465–97.
- 29. Comrey AL, Lee HB. A first course in factor analysis. 2nd ed. New York, NY: Psychology Press, Taylor & Francis Group; 2013.
- 30. World Health Organization Process of Translation and Adaptation of Instruments. [(accessed on 12 October 2020)]. Available online: http://www.who.in t/substance_abuse/research_tools/translation/en/
- Kalfoss M. Translation and adaptation of questionnaires: A nursing challenge. SAGE Open Nurs. 2019;5:2377960818816810. https://doi.org/10.1177/237796 0818816810.
- 32. Seyf AA. Measurement, test and educational evaluation. 7th ed. Tehran: Douran; 2016.
- Lawshe CH. A quantitative approach to content validity. Pers Psychol. 1975;28:563–75.
- Harerimana A, Mtshali NG. Using exploratory and confirmatory factor analysis to understand the role of technology in nursing education. Nurse Educ Today. 2020;92:104490. https://doi.org/10.1016/j.nedt.2020.104490.
- Schreiber J, Nora A, Stage F, Barlow L, King J. Confirmatory factor analyses and structural equations modeling: an introduction and review. J EdU Res. 2006;99(6).
- Norman GR, Sloan JA, Wyrwich KW. Interpretation of changes in healthrelated quality of life: the remarkable universality of half a standard deviation. Med Care. 2003;41(5):582–92. https://doi.org/10.1097/01.MLR.0000062554.74 615.4C.
- Nunnally J, Bernstein I. Psychometric theory 3rd edition. New York: MacGraw-Hill; In.; 1994.
- Yen M, Lo LH. Examining test-retest reliability: an intra-class correlation approach. Nurs Res. 2002;51(1):59–62.
- Gulledge CM, Smith DG, Ziedas A, Muh SJ, Moutzouros V, Makhni EC. Floor and ceiling effects, time to completion, and question burden of PROMIS CAT domains among shoulder and knee patients undergoing nonoperative and operative treatment. JB JS Open Access. 2019;4(4):e0015. https://doi.org/10.2 106/JBJS.OA.19.00015.
- Mashayekh-Amiri S, Jafarabadi MA, Hosseinzadeh M, Kanani ES, Mirghafourvand M. Measurement properties of the Persian version of the breast cancer perception scale (BCPS) according to the COSMIN checklist. BMC Cancer. 2024;24(1):743. https://doi.org/10.1186/s12885-024-12493-2.
- Geerinck A, Alekna V, Beaudart C, Bautmans I, Cooper C, De Souza Orlandi F, et al. Standard error of measurement and smallest detectable change of the sarcopenia quality of life (SarQoL) questionnaire: an analysis of subjects from 9 validation studies. PLoS ONE. 2019;14(4):e0216065. https://doi.org/10.1371/ journal.pone.0216065.
- 42. Shrivastava SR, Shrivastava PS, Ramasamy J. Role of self-care in management of diabetes mellitus. J Diabetes Metab Disord. 2013;12(1):14. https://doi.org/1 0.1186/2251-6581-12-14.
- Haron Z, Sutan R, Zakaria R, Abdullah Mahdy Z. Self-care educational guide for mothers with gestational diabetes mellitus: A systematic review on identifying self-care domains, approaches, and their effectiveness. Belitung Nurs J. 2023;9(1):6–16. https://doi.org/10.33546/bnj.2396.
- Karavasileiadou S, Almegwely W, Alanazi A, Alyami H, Chatzimichailidou S. Self-management and self-efficacy of women with gestational diabetes mellitus: a systematic review. Glob Health Action. 2022;15(1):2087298. https://doi. org/10.1080/16549716.2022.2087298.
- Assaf EA, Al Sabbah H, Momani A, Al-Amer R, Al-Sa'ad A, Ababneh G. Factors influencing gestational diabetes self-care among pregnant women in a Syrian refugee camp in Jordan. PLoS ONE. 2024;19(2):e0297051. https://doi.org/ 10.1371/journal.pone.0297051.
- Xu Y, Savage C, Toobert D, Wei Pan, Whitmer K. Adaptation and testing of instruments to measure diabetes self-management in people with type 2 diabetes in Mainland China. J Transcult Nurs. 2008;19(3):234–42. https://doi.or g/10.1177/1043659608319239.

- of diabetes self-care activities questionnaire: an Indonesian version. Indones Nurs J Educ Clin. 2019;4(1):25–36. 48. Vincent D, McEwen MM, Pasvogel A. The validity and reliability of a Spanish
- version of the summary of diabetes self-care activities questionnaire. Nurs Res. 2008;57(2):101–6. https://doi.org/10.1097/01.NNR.0000313484.18670.ab.
- Mogre V, Abanga ZO, Tzelepis F, Johnson NA, Paul C. Psychometric evaluation of the summary of diabetes self-care activities measure in Ghanaian adults living with type 2 diabetes. Diabetes Res Clin Pract. 2019;149:98–106. https:// doi.org/10.1016/j.diabres.2019.02.004.
- Kamradt M, Bozorgmehr K, Krisam J, Freund T, Kiel M, Qreini M, et al. Assessing self-management in patients with diabetes mellitus type 2 in Germany: validation of a German version of the summary of diabetes Self-Care activities measure (SDSCA-G). Health Qual Life Outcomes. 2014;12:185. https://doi. org/10.1186/s12955-014-0185-1.
- Jalaludin M, Fuziah M, Hong J, Mohamad Adam B, Jamaiyah H. Reliability and validity of the revised summary of diabetes Self-Care activities (SDSCA) for Malaysian children and adolescents. Malays Fam Physician. 2012;7(2–3):10–20.
- Ansari RM, Harris MF, Hosseinzadeh H, Zwar N. The summary of an Urdu version of diabetes Self-Care activities measure: psychometric evaluation and validation. J Prim Care Community Health. 2020;11:2150132720935292. https: //doi.org/10.1177/2150132720935292.
- Chang M, Roman-Lantzy C, O'Neil SH, Reid MW, Borchert MS. Validity and reliability of CVI range assessment for clinical research (CVI range-CR): a longitudinal cohort study. BMJ Open Ophthalmol. 2022;7(1):e001144.
- Shrestha N. Factor analysis as a tool for survey analysis. Am J Appl Math Stat. 2021;9(1):4–11. https://doi.org/10.12691/ajams-9-1-2.
- Choi EJ, Nam M, Kim SH, Park CG, Toobert DJ, Yoo JS, Chu SH. Psychometric properties of a Korean version of the summary of diabetes self-care activities measure. Int J Nurs Stud. 2011;48(3):333–7. https://doi.org/10.1016/j.ijnurstu.2 010.08.007.

- Rasmussen L, Poulsen CW, Kampmann U, Smedegaard SB, Ovesen PG, Fuglsang J. Diet and healthy lifestyle in the management of gestational diabetes mellitus. Nutrients. 2020;12(10):3050. https://doi.org/10.3390/nu12103050.
- Harrison AL, Taylor NF, Frawley HC, Shields N. A consumer cocreated infographic improves short-term knowledge about physical activity and selfefficacy to exercise in women with gestational diabetes mellitus: a randomized trial. J Physiother. 2020;66(4):243–8. https://doi.org/10.1016/j.jphys.2020. 09.010.
- Rose AJ, Richter EA. Skeletal muscle glucose uptake during exercise: how is it regulated? Physiol (Bethesda). 2005;20:260–70. https://doi.org/10.1152/physi ol.00012.2005.
- Symons Downs D, Ulbrecht JS. Understanding exercise beliefs and behaviors in women with gestational diabetes mellitus. Diabetes Care. 2006;29(2):236– 40. https://doi.org/10.2337/diacare.29.02.06.dc05-1262.
- Clar C, Barnard K, Cummins E, Royle P, Waugh N, Aberdeen Health Technology Assessment Group. Self-monitoring of blood glucose in type 2 diabetes: systematic review. Health Technol Assess. 2010;14(12):1–140. https://doi.org/ 10.3310/hta14120.
- Smyth S, Mulligan K, Rutter E, Harrington L, Hatunic M, Higgins MF. Attitudes of women with gestational diabetes toward diet and exercise: a qualitative study. J Matern Fetal Neonatal Med. 2023;36(1):2155045. https://doi.org/10.10 80/14767058.2022.2155045.
- 62. Maghalian M, Abbasalizadeh F, Mohammad-Alizadeh-Charandabi S, Ghanbari-Homaie S, Mirghafourvand M. Implementation and evaluation of the centering pregnancy group prenatal care model in pregnant women with diabetes: a convergent parallel mixed methods study protocol. Reprod Health. 2024;21(1):54.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.